SUICIDAL IDEATION						
Ask questions 1 and 2. If both are negative, proceed to "S	Suicidal Behavior" section. If the answer to		e: Time	n		
estion 2 is "yes", ask questions 3, 4 and 5. If the answer to question 1 and/or 2 is "yes", complete			e/They Most	Past 1 month		
"Intensity of Ideation" section below.			cidal	mor	uui	
1. Wish to be Dead		Jui	iuui			
Person endorses thoughts about a wish to be dead or not alive anymore	or wish to fall asleep and not wake up.	Yes	No	Yes	No	
Have you wished you were dead or wished you could go to sleep and n						
	-					
If yes, describe:						
2. Non-Specific Active Suicidal Thoughts		<b>X</b> 7		<b>X</b> 7	<b>N</b> 7	
General non-specific thoughts of wanting to end one's life/die by suicid		Yes	No	Yes	No	
ways to kill oneself/associated methods, intent, or plan during the assess Have you actually had any thoughts of killing yourself?	ment period.					
Have you actually had any thoughts of killing yourself?						
If yes, describe:						
2 A - 4						
3. Active Suicidal Ideation with Any Methods (Not Plan)		Yes	No	Yes	No	
Person endorses thoughts of suicide and has thought of at least one method during the assessment period. This is different than a specific plan with time, place or method details worked out (e.g., thought of method to kill self but not a specific plan). Includes person						
who would say, "I thought about taking an overdose but I never made a						
itand I would never go through with it."						
Have you been thinking about how you might do this?						
If yes, describe:						
4. Active Suicidal Ideation with Some Intent to Act, with	out Specific Plan					
4. Active Suicidal Ideation with Some Intent to Act, with Active suicidal thoughts of killing oneself and person reports having so		Yes	No	Yes	No	
thoughts but I definitely will not do anything about them."	ne ment to act on such moughts, as opposed to Thave the			_	_	
Have you had these thoughts and had some intention of acting on the	<i>n</i> ?					
If yes, describe:						
5. Active Suicidal Ideation with Specific Plan and Intent						
Thoughts of killing oneself with details of plan fully or partially worked		Yes	No	Yes	No	
Have you started to work out or worked out the details of how to kill you						
If yes, describe:						
INTENSITY OF IDEATION						
The following features should be rated with respect to the most.	severe type of ideation (i.e., 1-5 from above, with 1 being					
the least severe and 5 being the most severe). Ask about time he	/she/they were feeling the most suicidal.					
Lifetime - Most Severe Ideation:			ost	Mo		
<i>Type</i> # (1-5)	Description of Ideation	561	vere	Sev	ere	
Recent - Most Severe Ideation:						
<i>Type</i> # (1-5)	Description of Ideation					
Frequency						
How many times have you had these thoughts?						
(1) Less than once a week (2) Once a week (3) 2-5 times in week (4) Daily or almost daily (5) Many times each day						
Duration						
When you have the thoughts how long do they last?						
(1) Fleeting - few seconds or minutes	(4) 4-8 hours/most of day					
(2) Less than 1 hour/some of the time	(5) More than 8 hours/persistent or continuous					
(3) 1-4 hours/a lot of time						
Controllability	ing to dia if you want to?					
Could/can you stop thinking about killing yourself or want (1) Easily able to control thoughts	(4) Can control thoughts with a lot of difficulty					
(1) Easily able to control moughts (2) Can control thoughts with little difficulty	(4) Can control moughts with a lot of difficulty (5) Unable to control thoughts					
(3) Can control thoughts with some difficulty	(0) Does not attempt to control thoughts					
Deterrents						
Are there things - anyone or anything (e.g., family, religion	, pain of death) - that stopped you from wanting to					
die or acting on thoughts of suicide?						
(1) Deterrents definitely stopped you from attempting suicide	(4) Deterrents most likely did not stop you					
(2) Deterrents probably stopped you	(5) Deterrents definitely did not stop you					
(3) Uncertain that deterrents stopped you	(0) Does not apply					
Reasons for Ideation						
What sort of reasons did you have for thinking about want						
or stop the way you were feeling (in other words you could						
feeling) or was it to get attention, revenge or a reaction from						
<ul><li>(1) Completely to get attention, revenge or a reaction from others</li><li>(2) Mostly to get attention, revenge or a reaction from others</li></ul>	(4) Mostly to end or stop the pain (you couldn't go on living with the pain or how you were feeling)					
(2) Mostly to get attention, revenge or a reaction from others (3) Equally to get attention, revenge or a reaction from others	(5) Completely to end or stop the pain (you couldn't go on					
and to end/stop the pain	living with the pain or how you were feeling)					

SUICIDAL BEHAVIOR (Check all that apply, so long as these are separate events; must ask about all types)			Lifetime		Past 3 months	
Actual Attempt: A potentially self-injurious act undertaken with at least some wish to die, <i>as a result of act</i> . Behavior was in part thought of as method to kill oneself. Intent does not have to be 100%. If there is <i>any</i> intent/desire to die associated with the act, then it can be considered an actual suicide attempt. <i>There does not have to be any injury or harm</i> , just the potential for injury or harm. If person pulls trigger while gun is in mouth but gun is broken so no injury results, this is considered an attempt. Inferring Intent: Even if an individual denies intent/wish to die, it may be inferred clinically from the behavior or circumstances. For example, a highly lethal act that is clearly not an accident so no other intent but suicide can be inferred (e.g., gunshot to head, jumping from window of a high floor/story). Also, if someone denies intent to die, but they thought that what they did could be lethal, intent may be inferred. <i>Have you made a suicide attempt?</i>				Yes	No	
Have you made a succide altempt: Have you done anything to harm yourself? Have you done anything dangerous where you could have died? What did you do? Did you as a way to end your life? Did you want to die (even a little) when you? Were you trying to end your life when you? Or Did you think it was possible you could have died from? Or did you do it purely for other reasons / without ANY intention of killing yourself (like to relieve stress, feel better, get sympathy, or get something else to happen)? (Self-Injurious Behavior without suicidal intent)				Total # of Attempts		
If yes, describe: Has person engaged in Non-Suicidal Self-Injurious Behavior?			No	Yes	No	
<ul> <li>Interrupted Attempt:</li> <li>When person is interrupted (by an outside circumstance) from starting the potentially self-injurious act (<i>if not for that, actual attempt would have occurred</i>).</li> <li>Overdose: Person has pills in hand but is stopped from ingesting. Once they ingest any pills, this becomes an attempt rather than an interrupted attempt. Shooting: Person has gun pointed toward self, gun is taken away by someone else, or is somehow prevented from pulling trigger. Once they pull the trigger, even if the gun fails to fire, it is an attempt. Jumping: Person is poised to jump, is grabbed and taken down from ledge. Hanging: Person has noose around neck but has not yet started to hang - is stopped from doing so.</li> <li>Has there been a time when you started to do something to end your life but someone or something stopped you before you actually did anything?</li> <li>If yes, describe:</li> </ul>				Yes No		
				Total # of interrupted		
Aborted or Self-Interrupted Attempt: When person begins to take steps toward making a suicide attempt but stops themselves before they actually have engaged in any self- destructive behavior. Examples are similar to interrupted attempts, except that the individual stops him/herself, instead of being stopped by something else. Has there been a time when you started to do something to try to end your life but you stopped yourself before you actually did anything? If yes, describe:			No	f Total # of aborted or self-		
<b>Preparatory Acts or Behavior:</b> Acts or preparation towards imminently making a suicide attempt. This can include anything beyond a verbalization or thought, such as assembling a specific method (e.g., buying pills, purchasing a gun) or preparing for one's death by suicide (e.g., giving things away, writing a suicide note).			No	Yes	No	
Have you taken any steps towards making a suicide attempt or preparing to kill yourself (such as collecting pills, getting a gun, giving valuables away or writing a suicide note)? If yes, describe:			Total # of preparatory acts		Total # of preparatory acts	
	Attempt	Most Leth Attempt Date:		Initial/Fin Attempt Date:	rst	
<ul> <li>Actual Lethality/Medical Damage:</li> <li>0. No physical damage or very minor physical damage (e.g., surface scratches).</li> <li>1. Minor physical damage (e.g., lethargic speech; first-degree burns; mild bleeding; sprains).</li> <li>2. Moderate physical damage; medical attention needed (e.g., conscious but sleepy, somewhat responsive; second-degree burns; bleeding of major vessel).</li> <li>3. Moderately severe physical damage; medical hospitalization and likely intensive care required (e.g., comatose with reflexes intact; third-degree burns less than 20% of body; extensive blood loss but can recover; major fractures).</li> <li>4. Severe physical damage; medical hospitalization with intensive care required (e.g., comatose without reflexes; third-degree burns over 20% of body; extensive blood loss with unstable vital signs; major damage to a vital area).</li> <li>5. Death</li> </ul>	Enter Code	Enter C	'ode	Enter (	Code	
Potential Lethality: Only Answer if Actual Lethality=0       Enter Code         Likely lethality of actual attempt if no medical damage (the following examples, while having no actual medical damage, had potential for very serious lethality: put gun in mouth and pulled the trigger but gun fails to fire so no medical damage; laying on train tracks with oncoming train but pulled away before run over).		Enter Code		Enter Code		
0 = Behavior not likely to result in injury 1 = Behavior likely to result in injury but not likely to cause death 2 = Behavior likely to result in death despite available medical care						